PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: A61K 39/12, C07H 21/02, C12N 7/04, 15/40, 15/86

(11) International Publication Number:

WO 96/37220

A1

(43) International Publication Date:

28 November 1996 (28.11.96)

(21) International Application Number:

PCT/US96/07457

(22) International Filing Date:

22 May 1996 (22.05.96)

(30) Priority Data:

08/446,932

23 May 1995 (23.05.95)

US

(60) Parent Application or Grant

(63) Related by Continuation

US Filed on 08/446,932 (CIP)

23 May 1995 (23.05.95)

(71) Applicant (for all designated States except US): THE UNI-VERSITY OF NORTH CAROLINA AT CHAPEL HILL [US/US]; 300 Bynum Hall, Chapel Hill, NC 27599 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JOHNSTON, Robert, E. [US/US]; 101 Marin Place, Chapel Hill, NC 27516 (US). DAVIS, Nancy, L. [US/US]; 415 Sharon Road, Chapel Hill, NC 27514 (US). SIMPSON, Dennis, A. [US/US]; Apartment 50, 400 Davie Road, Carrboro, NC 27510 (US).

(74) Agents: SIBLEY, Kenneth, D. et al.; Bell, Seltzer, Park & Gibson, P.O. Drawer 34009, Charlotte, NC 28234 (US).

(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: cDNA CLONE FOR SOUTH AFRICAN ARBOVIRUS NO. 86

(57) Abstract

The present invention provides a recombinant DNA comprising a cDNA coding for an infectious South African Arbovirus No. 86 (S.A.AR86) virus RNA transcript and a heterologous promoter positioned upstream from the cDNA and operatively associated therewith. The present invention also provides an infectious RNA transcript encoded by the cDNA, and infectious attenuated viral particles produced from cells transfected with the RNA transcript encoded by the cDNA.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	4		·		
AT	Armenia	GB	United Kingdom	MW	Malawi
ΑŬ	Austria	GE	Georgia	MX	Mexico
BB	Australia	GN	Guinea	NE	Niger
BE	Barbados	GR	Greece	NL	Netherlands
	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic		of Korea	SE	Sweden .
CG	Congo	KR	Republic of Korea	SG	Singapore
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	
CN .	China	LR	Liberia	SZ	Senegal
CS	Czechoslovakia .	LT	Lithuania		Swaziland
CZ	Czech Republic	LU	Luxembourg	TD	Chad
DE	Germany	LV	Latvia	TG	Togo
DK	Denmark	MC	Моласо	TJ	Tajikistan
EE	Estonia	MD	Republic of Moldova	TT	Trinidad and Tobago
ES	Spain	MG	Madagascar	UA	Ukraine
FI	Finland	ML	Mali	UG	Uganda
FR	France	MN	Mongolia	US	United States of America
GA ·	Gabon	MR	Mauritania	UZ	Uzbekistan
		****	trames and tig	VN	Viet Nam

20

25

cDNA CLONE FOR SOUTH AFRICAN ARBOVIRUS NO. 86 Field of the Invention

The present invention relates to live attenuated vaccines in general, and particularly relates to attenuated vaccines produced from South African Arbovirus No. 86 (S.A.AR86) virus.

Background of the Invention

This invention was made with government support under Grant No. 2-ROIA122186 (09-13) awarded by the National Institutes of Health. The government has certain rights in the invention.

South African Arbovirus No. 86 (S.A.AR86) is an isolate of Sindbis 10 virus. S.A.AR 86 was originally isolated from mosquitoes. The virus is antigenically related to Sindbis virus and to two other Sindbis virus isolates, Girdwood S.A. and Ockelbo82. See, Malherbe et al., South African Medical Journal, 37:547 (1963) and Niklasson et al., Am. J. Trop. Med. Hyg. 33:1212 (1984), respectively. The latter is associated with a human disease, also known 15 as Ockelbo. Sindbis virus is the prototype member of the alphavirus genus of the The Sindbis virus includes various strains, including family Togaviridae. S.A.AR86 and Sindbis AR339. The genome of Sindbis viruses consists of a single strand of RNA which contains the information for the viral genes, and which is infectious when introduced into the cytoplasm of cells.

Full-length cDNA clones of positive-strand RNA viruses are important tools for the study of the biology of viruses including Sindbis viruses. It is known with respect to several viral systems that *in vitro* transcripts of cDNA clones, and in some cases the cDNA itself, can initiate a complete and productive infectious cycle upon introduction into susceptible cells. *See* Racaniello et al., *Science* 214:916 (1981); Ahlquist et al., *Proc. Natl. Acad. Sci. USA* 81:7066 (1984); Kaplan et al., *Proc. Natl. Acad. Sci. USA* 82:8424 (1985); Mizutani et al., *J. Virol.* 56:628 (1985); van der Werf, *Proc. Natl. Acad. Sci. USA* 83:2330 (1986); Rice et al., *J. Virol.* 61:3809 (1987); and Vos et al., *Virology* 165:33 (1988). This has made it possible to test progeny virus for phenotypic

10

15

20

25

manifestations of directed mutations and recombinations which have been introduced into the cDNA clone. Pathogenesis studies with several positive-strand viruses, including the picornaviruses and the alphaviruses have been advanced significantly by the use of full-length cDNA clones.

As another useful application, live attenuated viral vaccines may be produced using full-length cDNA clones. Live attenuated viral vaccines are among the most successful means of controlling viral disease. However for some virus pathogens, immunization with a live virus strain may be either impractical or unsafe. Sindbis virus is subclinical in humans, but is closely related to other viruses which do induce clinical diseases in humans, such as Ockelbo, an epidemic polyarthritis syndrome common in areas of Scandinavia and Northern Europe. Accordingly, Sindbis virus vaccines are desirable for producing an immunogenic response to such clinical diseases. Sindbis virus vaccines are also desirable as viral carriers in virus constructs which express genes encoding immunizing antigens for other viruses. See U.S. Patent No. 5,217,879 to Huang et al. Huang et al. describes Sindbis infectious viral vectors. However, the reference does not describe the cDNA sequence of S.A.AR86 virus, or clones or viral vectors produced therefrom.

Accordingly, there remains a need in the art for full-length cDNA clones of positive-strand RNA viruses, such as the S.A.AR86 strain of Sindbis. In addition, there is a need in the art for full-length cDNA clones of S.A.AR86 encoding infectious RNA transcripts. Further, there remains a need in the art for cDNA clones of S.A.AR86 which encode RNA transcripts which may be used to produce infectious attenuated viral particles, and methods of producing such viral particles.

Summary of the Invention

As a first aspect, the present invention provides a recombinant DNA comprising a cDNA coding for an infectious South African Arbovirus No. 86 (S.A.AR86) virus RNA transcript and a heterologous promoter positioned upstream from the cDNA and operatively associated therewith. The cDNA is selected from the group consisting of (i) cDNA having the sequence given herein

WO 96/37220

15

20

25

30

as SEQ ID NO.: 1, (ii) cDNA having the same RNA coding sequence as the cDNA given herein as SEQ ID NO.: 1, and (iii) cDNA according to (i) or (ii) above and further containing at least one attenuating mutation. Preferably at least one attenuating mutation is included in the cDNA, and more preferably at least two attenuating mutations are included in the cDNA. Attenuating mutations may, for example, be provided in any of the nsP1, E2, and nsP2 coding regions of the cDNA. Preferably at least one silent mutation is included in the cDNA in addition to the attenuating mutation(s).

As a second aspect, the present invention provides an infectious RNA transcript encoded by the cDNA.

As a third aspect, the present invention provides infectious attenuated viral particles containing the RNA transcript encoded by the cDNA.

The foregoing and other aspects of the present invention are explained in detail in the detailed description set forth below.

Brief Description of the Drawings

Figure 1 shows the relationship of the 3' half of the nsP3 gene among various Sindbis-like isolates.

Figure 2 shows the replacement of the AR339-derived cDNAs into the plasmid pToto1101 background.

Detailed Description of the Invention

The South African Arbovirus No. 86 (S.A.AR86) viral clones employed in practicing the present invention are genomic clones which code for an RNA transcript, which RNA transcript is capable of producing live encapsidated S.A.AR86 virus when used to transfect a S.A.AR86 virus-permissive cell.

S.A.AR86 virus-permissive cells are cells which, upon transfection with the viral RNA transcript, are capable of producing viral particles. The S.A.AR86 virus has a broad host cell range. Examples of suitable host cells include, but are not limited to Vero cells, baby hamster kidney (BHK-21) cells, and chicken embryo fibroblast cells. Uptake of the RNA into the cells can be

10

15

20

25

30

achieved by any suitable means, such as for example, by treating the cells with DEAE-dextran, treating the cells with "LIPOFECTIN^{RM}", and by electrophoresis, with electrophoresis being the currently preferred means of achieving RNA uptake into the host cells.

The phrases "attenuating mutation" and "attenuating amino acid," as used herein, mean a nucleotide mutation or an amino acid coded for in view of such a mutation which result in a decreased probability of causing disease in its host (i.e., a loss of virulence), in accordance with standard terminology in the art. See, e.g., B. Davis, et al., Microbiology 132 (3d ed. 1980), whether the mutation be a substitution mutation or an in-frame deletion mutation. The phrase "attenuating mutation" excludes mutations which would be lethal to the virus.

The phrase "silent mutation" as used herein refers to mutations in the cDNA coding sequence which do not produce mutations in the corresponding protein sequence translated therefrom.

The cDNA clone has a sequence as given herein as SEQ ID NO.: 1. Alternatively, the cDNA clone may have a sequence which differs from the cDNA of SEO ID NO.: 1, but which has the same RNA coding sequence as the cDNA given herein as SEQ ID NO.: 1. Thus, the cDNA clone may include one or more silent mutations. For example, the clone sequence may differ from the wild-type S.A.AR86 sequence given herein as SEQ ID NO.: 1, by the inclusion of silent mutations at any or all of nucleotides 215, 3863, 5984, and 9113. The silent mutations at the foregoing nucleotides may be substitution or in-frame deletion mutations, such as for example, the substitution of guanine for adenine at nucleotide 215 of the cDNA sequence given herein as SEQ ID NO.: 1; or the substitution of guanine for cytosine at nucleotide 3863 of the cDNA sequence given herein as SEQ ID NO.: 1; or the substitution of guanine for adenine at nucleotide 5984 of the cDNA sequence given herein as SEQ ID NO.: 1; or the substitution of cytosine for thymine at nucleotide 9113 of the cDNA sequence given herein as SEQ ID NO.: 1. In yet another emobidment, the cDNA clone has a sequence according to either of the foregoing described sequences, but which also includes attenuating mutations. The attenuating mutations being described more fully hereinafter.

15

20

25

30

Promoter sequences and S.A.AR86 virus cDNA clones are operatively associated in the present invention such that the promoter causes the cDNA clone to be transcribed in the presence of an RNA polymerase which binds to the promoter. The promoter is positioned on the 5' end (with respect to the virion RNA sequence), or "upstream" from, the cDNA clone. An excessive number of nucleotides between the promoter sequence and the cDNA clone will result in the inoperability of the construct. Hence, the number of nucleotides between the promoter seuqence and the cDNA clone is preferably not more than eight, more preferably not more than five, still more preferably not more than three, and most preferably not more than one. Examples of promoters which are useful in the cDNA sequences of the present invention include, but are not limited to T3 promoters, T7 promoters, and SP6 promoters. The DNA sequence of the present invention may reside in any suitable transcription vector. The DNA sequence preferably has a complementary DNA sequence bonded thereto so that the double-stranded sequence will serve as an active template for RNA polymerase. The transcription vector preferably comprises a plasmid. When the DNA sequence comprises a plasmid, it is preferred that a unique restriction site be provided 3' (with respect to the virion RNA sequence) to (i.e., "downstream" from) the cDNA clone. This provides a means for linearizing the DNA sequence to allow the transcription of genome-length RNA in vitro.

The cDNA clone can be generated by any of a variety of suitable methods known to those skilled in the art. A preferred method is the method set forth in U.S. Patent No. 5,185,440 to Davis et al., and Gubler et al., Gene 25:263 (1983), the disclosures of which are incorporated herein by reference in their entirety. Attenuating mutations of S.A.AR86 are identified by sequencing attenuated strains of the S.A.AR86 virus and comparing the sequence of the attenuated strain with the sequence of the corresponding wild-type virus. Serial passage techniques for the generation of attenuated strains may be carried out in accordance with known procedures. Preferably, the atenuated strains are generated by selecting strains at each passage during serial passage in cell culture which either grow rapidly or penetrate the cell more rapidly. This selection process, which reduces the number of serial passages required to obtain attenuated strains.

15

20

25

30

is known. See, e.g., Olmstead et al., Science 225:424 (1984); and Johnston et al., Virology 162:437 (1988). the disclosures of which are incorporated herein by reference in their entirety. cDNA clones may be modified to incorporate attenuating mutations by site-directed mutagenesis in accordance with known procedures. An exemplary technique is that of Kunkel Proc. Natl. Acad. Sci. (USA) 82:488 (1985). These same techniques may be used to join the heterologous promoter to the cDNA clone.

RNA is preferably synthesized from the DNA sequence in vitro using purified RNA polymerase in the presence of ribonucleotide triphosphates in accordance with conventional techniques.

Pharmaceutical compositions, such as vaccines, containing the S.A.AR86 clone of the present invention comprise an immunogenic amount of a live attenutated virus as disclosed herein in combination with a pharmaceutically acceptable carrier. An "effective immunogenic amount" is an amount of the attenuated virus sufficient to evoke an immune response in the subject to which the vaccine is administered. An amount of about 10¹ to about 10⁵ plaque forming units per dose is believed to be suitable, depending upon the age and species of the subject being treated. Examples of suitable pharmaceutically acceptable carriers include, but are not limited to, sterile pyrogen-free water and sterile pyrogen-free physiological saline solution. Subjects which may be administered immunogenic amounts of the live attenuated viruses of the present invention include both human and animal (e.g., horse, donkey, mouse, hamster, or monkey) subjects. Administration may be by a suitable means, such as intraperitoneoal, intracerebral or intramuscular injection.

Complimentary DNA clones of the S.A.AR86 virus are made in accordance with the procedures described herein, as supplemented with procedures known in the art. We employed as a starting material, the S.A.AR86 virus.

A first exemplary attenuating substitution mutation in a S.A.AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably isoleucine, at nsP1 amino acid residue 538.

15

25

30

A second exemplary attenuating substitution mutation in a S.A. AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably threonine, at E2 amino acid residue 304.

A third exemplary attenuating substitution mutation in a S.A.AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably lysine, at E2 amino acid residue 314.

A fourth exemplary attenuating substitution mutation in a S.A.AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably valine, at E2 amino acid residue 372.

A fifth exemplary attenuating substitution mutation in a S.A.AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably alanine, at E2 amino acid residue 376.

One embodiment of the present invention contains, in combination, said attenuating substitution mutations at E2 amino acid residues 304, 314, 372, and 376.

A sixth exemplary attenuating substitution mutation in a S.A.AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably glycine, at nsP2 amino acid residue 96.

A seventh exemplary attenuating substitution mutation in a S.A.AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably valine, at nsP2 amino acid residue 372.

One embodiment of the present invention contains, in combination, said attenuating substitution mutations at nsP2 amino acid residues 96 and 372.

An eighth exemplary attenuating substitution mutation in an S.A.AR86 clone useful in practicing the present invention is a substitution

15

20

25

30

mutation which codes for an attenuating amino acid, preferably leucine, at nsP2 amino acid residue 529.

A nineth exemplary attenuating substitution mutation in an S.A.AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably asparagine, at nsP2 amino acid residue 571.

A tenth exemplary attenuating substitution mutation in an S.A.AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably arginine, at nsP2 amino acid residue 682.

An eleventh exemplary attenuating substitution mutation in an S.A.AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably arginine, at nsP2 amino acid residue 804.

A twelveth exemplary attenuating substitution mutation in an S.A.AR86 clone useful in practicing the present invention is a substitution mutation which codes for an attenuating amino acid, preferably arginine, at nsP3 amino acid residue 22.

One embodiment of the present invention contains, in combination, said attenuating substitution mutations at nsP2 amino acid residues 529, 571, 682, and 804, and at nsP3 amino acid residue 22.

The cDNA clones according to the present invention are useful for the preparation of pharmaceutical formulations, such as vaccines, as described above. In addition, the cDNA clones of the present invention are useful for administration to animals for the purpose of producing antibodies to the S.A.AR86 virus, which antibodies may be collected and used in known diagnostic techniques for the detection of S.A.AR86 virus.

The following examples are provided to illustrate the present invention, and should not be construed as limiting thereof. In these examples, nt means nucleotide.

EXAMPLE 1

Relationship of S.A.AR86 Clone to Other Sindbis Strains

At the nucleotide level, S.A.AR86 differs from the consensus sequence of Sindbis strain AR339 by 685 nucleotides (111 amino acids). From the published Sindbis HR sequence (See, Strauss et al., Virol. 133:92 (1984)), S.A.AR86 differs by 704 nucleotides and 119 amino acids. S.A.AR86 differs from the sequence of Ockelbo82 (See, Shirako et al., Virol. 182:753 (1991)) by 430 nucleotides (67 amino acids). Included in these differences are several insertions and deletions present in S.A.AR86 relative to the Sindbis sequences The relationship of the 3' half of the nsP3 gene among various Sindbis-like isolates is shown in Figure 1.

EXAMPLE 2

Observed Mortality in Mice Infected with S.A.AR86

When adult mice are inoculated intracerebrally (i.c.) with low doses of S.A.AR86, 100% motality is observed. The methods employed for evaluating the instance mortality in mice are set forth in D. Russell, et al., J. Virol. 63(4):1619 (1989), the disclosure of which is incorporated herein by reference in its entirety. This high mortality rate is unique among "Sindbis-like" viruses which are characteristically avirulent in adult mice. Also unique in the S.A.AR86 strain is the cys substitution for the opal stop codon normally found between the nsP3 and nsP4 genes.

EXAMPLE 3

Construction of S.A.AR86 Clone

The S.A.AR86 clone is constructed by substituting partial cDNA clones of S.A.AR86 genomic RNA into pTR5000, one of a series of Sindbis AR339 cDNA clones. Construction of pTR5000 (a full-length cDNA clone of Sindbis following the SP6 phage promoter and containing mostly Sindbis AR339 sequences) is accomplished by sequential replacement of AR339-derived cDNAs into the plasmid pToto1101 background, according to the technique described in

Rice et al., J. Virol. 61:3809 (1987). The replacement of the AR339-derived cDNAs into the plasmid pToto1101 background is shown in Figure 2.

Production of the cDNAs used in constructing pTR5000 has been described previously in Polo et al., J. Virol. 62:2124 (1988), as has the construction of pTR2000, see Polo et al., J. Virol. 62:2124 (1988) and Polo et al., J. Virol. 64:4438 (1990). Nucleotide numbering follows that of Strauss et al. Virol. 133:92 (1984).

The Stul (nt8571) to SacII (nt11484) fragment of pToto1101 is removed and replaced with the analogous fragment from clone pSB4 to form pTR2000, using the loss of the pToto1101 StuI site at nt10770 (a site not present in AR339) as a screen, as shown in Figure 2. The sequences of AR339 and pToto1101 are identical from nt11485 to the 3'-end (nt11703). Therefore, as shown in Figure 2, these 3' sequences are of AR339 origin. Construction of pTR3000 is accomplished by replacement of the BssHII (nt9804) to StuI (nt8571) fragment of pSB3 into pTR2000 from which the analogous fragment had been removed. The AfIII site found at nt8835 in pToto1101 but absent in AR339 is used to screen the recombinants. An AR339 fragment from pSB1, SpeI (nt5262) to BssHII, is used to replace the SpeI-BssHII fragment from pTR2000, using the AfIII screen and forming pTR4000. To construct pTR5000, pSB5 is subcloned into pUC119, and the PstI site at nt3953 is ablated using site-directed mutagenesis, as described in Kunkel, Methods Enzymol. 154:367 (1987), to change nt3950 from U to C. The ClaI (nt2713) to SpeI fragment is removed from the mutagenized subclone and for pTR5000, is used to replace the analogous fragment of pTR4000 using the ablated PstI site as a screen.

15

20

25

30

Partial cDNA clones of S.A.AR86 were obtained using classical reverse transcriptase (RT) procedures according to Polo et al., J. Virol. 62:2124 (1988), as well as RT-PCR protocols according to Heidner et al., J. Virol. 68:2683 (1994). These cDNA clones are used to replace analogous portions of the clone pTR5000, culminating in the construction of a full-length cDNA of the S.A.AR86 genomic sequence downstream of an SP6 promoter and followed by a poly (A) tract and a unique XbaI site. During the course of replacing S.A.AR86 sequences into pTR5000, it was observed that the pTR5000 nonstructural proteins

are incompatible with those of S.A.AR86, so that the chimeric clones yielded transcripts which are not infectious for baby hamster kidney (BHK-21) cells. In addition, the restriction sites in the nonstructural region of S.A.AR86 are very different from the Sindbis AR339-based clones. The first complete S.A.AR86 clone, pS10, also failed to yield infectious transcripts.

The construction is repeated, beginning with pTR5000, using sequences derived from the same partial cDNA clones of S.A.AR86 as are used for the construction of pS10 except for nucleotides 3171 to 6410 (numbering from nucleotide 1 of the S.A.AR86 sequence), which are derived by RT-PCR of the genomic RNA. The resulting construct, pS22, gives infectious transcripts, but the virus derived therefrom is temperature-sensitive. Replacement of nucleotides 3171 to 6410 with an alternative RT-PCR derived cDNA corrected the temperature-sensitive defect, yielding clone pS24.

10

15

20

25

30

EXAMPLE 4

Observed Mortality in Mice Infected with S.A.AR86

Upon i.c. inoculation virus derived from pS24 is avirulent, whereas S.A.AR86 caused 100% mortality. Clearly, pS24 contained one or more mutations which were strongly attenuating. The complete sequence of pS24 was determined directly from pS24 and related clones. Comparison with the S.A.AR86 genomic RNA sequence reveals 5 mutations or clusters of mutations which are potentially associated with the avirulent phenotype of virus from pS24. These included the mutations or clusters of mutations indicated in clones pS56, pS51, and pS57, a mutation at ntl278 A-C (nsP1 407 K-Q), and a mutation at nt5972 T \rightarrow G (nsP3 228 N \rightarrow S). While said nsP1 407 substitution alone is a lethal mutation, the viruses are viable when the mutation co-exists with a serine at the nsP3 amino acid residue 228. Clone pS24 is corrected by by a combination of sitedirected mutagenesis and replacement of specific pS24 sequences with cDNA fragments which do not contain the subject mutations. The resulting cDNA is pS55 which contains 5' and 3' untranslated sequences identical to S.A.AR86 genomic RNA, no coding differences with S.A.AR86 genomic RNA, and the 4

non-coding changes at nt 215 A-G, nt 3863 C-G, nt 5984 A-G, and nt 9113 T-C. Virus derived from pS55 is indistinguishable from native S.A.AR86 in tests of virulence, and growth in adult mice and by histopathological analysis of tissues from infected animals. These results, which are reported in Table 1 below, indicate that virus derived from clone pS55 accurately reflects native S.A.AR86 both in terms of coding sequence and *in vivo* phenotype.

Table 1
VIRULENCE IN MICE

	AGE OF MICE	S.A.AR86	\$55	S56 ²	S51 ³
10	4 to 6 Weeks				
•	Mortality	100%	100%	80%	20%
	AST ¹	6.36 ± 1.39	6.37 ± 1.4	9.37±1.77	8.5 ± 0.7

Average Survival Time in days.

20

25

The 3 mutations or clusters of mutations in clones pS56, pS51, and pS57 are placed independently into the pS55 background. Virus derived from each of these clones is highly attenuated in adult mice inoculated i.c. In clone pS53, the mutations at nucleotides 6 and 1672 are combined, and the resulting virus is avirulent in the adult mouse model. The mutations in pS61 are present in pS48, an intermediate clone constructed during the repair of pS24. The virus from pS48 produced small plaques on BHK-21 cells. When these mutations are placed in the pS55 background, they also gave a highly attenuated phenotype (S61, virus derived from pS61, gave 33.3% mortality (6±4.2)).

The foregoing is illustrative of the present invention and is not to be construed as limiting thereof. The invention is defined by the following claims, with equivalents of the claims to be included therein.

²Virus isogenic with S55 except at nucleotide 6 (C→A).

³Virus isogenic with S55 except at nucleotide 1648 (C→T). This is nsP1 amino acid 538 (T→T). The isoleucine is the amino acid found in all other Sindbis isolates sequenced to date.

SEQUENCE LISTING

(1)	GENERAL	INFORMATION:

(i) APPLICANT: Johnston, Robert E. Simpson, Dennis Davis, Nancy L.

- (ii) TITLE OF INVENTION: cDNA Clone for South African Arbovirus No. 86
- (iii) NUMBER OF SEQUENCES: 1

(iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Kenneth D. Sibley

(B) STREET: Post Office Drawer 34009

(C) CITY: Charlotte (D) STATE: NC

(E) COUNTRY: USA

(F) ZIP: 28234

- (v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 (B) FILING DATE:
 (C) CLASSIFICATION:

- (viii) ATTORNEY/AGENT INFORMATION:
 (A) NAME: Sibley, Kenneth D.
 (B) REGISTRATION NUMBER: 31,665
 (C) REFERENCE/DOCKET NUMBER: 5470-118
 - (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (919) 881-3140 (B) TELEFAX: (919) 881-3175 (C) TELEX: 575102
- (2) INFORMATION FOR SEQ ID NO:1:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11663 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: cDNA
 - (iii) HYPOTHETICAL: NO
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATTGGCGGCG TAGTACACAC TATTGAATCA AACAGCCGAC CAATTGCACT ACCATCACAA 60

TGGAGAAGCC AGTAGTTAAC GTAGACGTAG ACCCTCAGAG TCCGTTTGTC GTGCAACTGC

120

AAAAGAGCTT CCCGCAATTT GAGGTAGTAG CACAGCAGGT CACTCCAAAT GACCATGCTA

ATGCCAGAGC	ATTTTCGCAT	CTGGCCAGTA	AACTAATCGA	GCTGGAGGTT	CCTACCACAG	240
CGACGATTTT	GGAÇATAGGC	AGCGCACCGG	CTCGTAGAAT	GTTTTCCGAG	CACCAGTACC	300
ATTGCGTTTG	CCCCATGCGT	AGTCCAGAAG	ACCCGGACCG	CATGATGAAA	TATGCCAGCA	360
AACTGGCGGA	AAAAGCATGT	AAGATTACAA	ACAAGAACTT	GCATGAGAAG	ATCAAGGACC	420
TCCGGACCGT	ACTTGATACA	CCGGATGCTG	AAACGCCATC	ACTCTGCTTC	CACAACGATG	480
TTACCTGCAA	CACGCGTGCC	GAGTACTCCG	TCATGCAGGA	CGTGTACATC	AACGCTCCCG	540
GAACTATTTA	CCACCAGGCT	ATGAAAGGCG	TGCGGACCCT	GTACTGGATT	GGCTTCGACA	600
CCACCCAGTT	CATGTTCTCG	GCTATGGCAG	GTTCGTACCC	TGCATACAAC	ACCAACTGGG	660
CCGACGAAAA	AGTCCTTGAA	GCGCGTAACA	TCGGACTCTG	CÁGCACAAAG	CTGAGTGAAG	720
GCAGGACAGG	AAAGTTGTCG	ATAATGAGGA	AGAAGGAGTT	GAAGCCCGGG	TCACGGGTTT	780
ATTTCTCCGT	TGGATCGACA	CTTTACCCAG	AACACAGAGC	CAGCTTGCAG	AGCTGGCATC	840
TTCCATCGGT	GTTCCACTTG	AAAGGAAAGC	AGTCGTACAC	TTGCCGCTGT	GATACAGTGG	900
TGAGCTGCGA	AGGCTACGTA	GTGAAGAAAA	TCACCATCAG	TCCCGGGATC	ACGGGAGAAA	960
CCGTGGGATA	CGCGGTTACA	AACAATAGCG	AGGGCTTCTT	GCTATGCAAA	GTTACCGATA	1020
CAGTAAAAGG	AGAACGGGTA	TCGTTCCCCG	TGTGCACGTA	TATCCCGGCC	ACCATATGCG	1080
ATCAGATGAC	CGGCATAATG	GCCACGGATA	TCTCACCTGA	CGATGCACAA	AAACTTCTGG	1140
	•			CAGGAACACC		1200
			·	ATGGGCCAAG		1260
				CAAGCTTACA		1320
				CCCACCTGGA		1380
				ATCCGTATGG		1440
				ACCAAAGAAG		1500
•				TGCTTTCGAG		1560
		•		ATTAGTGGCA		1620
					ACCGGAGCAG	1680
					GACCGTATGA	1740
· ·					CTCGCACCAG	1800
					GGAAGGTATG	1860
· •					CCATGGCCAG	1920
					TTTGTGAACC	1980
					GAGCAGTACA	
					AAGAAGCGAT	
					AACCCGCCCT	
ATCACGAACT	AGCTCTTGAG	GGACTGAAGA	CTCGACCCGC	GGTCCCGTAC	AAGGTTGAAA	2220

CAATAGGAG	T GATAGGCACA	CCAGGATCGG	GCAAGTCAGC	TATCATCAAG	TCAACTGTCA	2280
CGGCACGTG	A TOTTGTTACC	AGCGGAAAGA	AAGAAAACTG	CCGCGAAATT	GAGGCCGACG	2340
TGCTACGGC	T GAGGGGCATG	CAGATCACGT	CGAAGACAGT	GGATTCGGTT	ATGCTCAACG	2400
GATGCCACA	A AGCCGTAGAA	GTGCTGTATG	TTGACGAAGC	GTTCCGGTGC	CACGCAGGAG	2460
CACTACTTG	C CTTGATTGCA	ATCGTCAGAC	CCCGTAAGAA	GGTAGTACTA	TGCGGAGACC	2520
CTAAGCAAT	G CGGATTCTTC	AACATGATGC	AACTAAAGGT	ACATTTCAAC	CACCCTGAAA	2580
AAGACATAT	g taccaagaca	TTCTACAAGT	TTATCTCCCG	ACGTTGCACA	CÁGCCAGTCA	2640
CGGCTATTG	T ATCGACACTG	CATTACGATG	GAAAAATGAA	AACCACAAAC	CCGTGCAAGA	2700
AGAACATCG	A AATCGACATT	ACAGGGCCA	CGAAGCCGAA	GCCAGGGGAC	ATCATCCTGA	2760
CATGTTTCC	G CGGGTGGGTT	AAGCAACTGC	AAATCGACTA	TCCCGGACAT	GAGGTAATGA	2820
CAGCCGCGG	C CTCACAAGGG	CTAACCAGAA	AAGGAGTATA	TGCCGTCCGG	CAAAAAGTCA	2880
ATGAAAACC	C GCTGTACGCG	ATCACATCAG	AGCATGTGAA	CGTGTTGCTC	ACCCGCACTG	2940
AGGACAGGC	T AGTATGGAAA	ACTTTACAGG	GCGACCCATG	GATTAAGCAG	CTCACTAACG	3000
TACCTAAAG	G AAATTITCAG	GCCACCATCG	AGGACTGGGA	AGCTGAACAC	AAGGGAATAA	3060
TTGCTGCGA	T AAACAGTCCC	GCTCCCCGTA	CCAATCCGTT	CAGCTGCAAG	ACTAACGTTT	3120
GCTGGGCGA	A AGCACTGGAA	CCGATACTGG	CCACGGCCGG	TATCGTACTT	ACCGGTTGCC	3180
AGTGGAGCG	A GCTGTTCCCA	CAGTTTGCGG	ATGACAAACC	ACACTCGGCC	ATCTACGCCT	3240
TAGACGTAA	T TTGCATTAAG	TTTTCGGCA	TGGACTTGAC	AAGCGGGCTG	TTTTCCAAAC	3300
AGAGCATCC	C GTTAACGTAC	CATCCTGCCG	ACTCAGCGAG	GCCAGTAGCT	CATTGGGACA	3360
ACAGCCCAG	G AACACGCAAG	TATGGGTACG	ATCACGCCGT	TGCCGCCGAA	CTCTCCCGTA	3420
GATTTCCGG	T GTTCCAGCTA	GCTGGGAAAG	GCACACAGCT	TGATTTGCAG	ACGGGCAGAA	3480
CTAGAGTTA	T CTCTGCACAG	CATAACTTGG	TCCCAGTGAA	CCGCAATCTC	CCTCACGCCT	3540
TAGTCCCCG	A GCACAAGGAG	AAACAACCCG	GCCCGGTCGA	AAAATTCTTG	AGCCAGTTCA	3600
AACACCACT	C CGTACTTGTG	ATCTCAGAGA	AAAAAATTGA	AGCTCCCCAC	AAGAGAATCG	3660
AATGGATCG	C CCCGATTGGC	ATAGCCGGCG	CAGATAAGAA	CTACAACCTG	GCTTTCGGGT	3720
TTCCGCCGC	A GGCACGGTAC	GACCTGGTGT	TCATCAATAT	TGGAACTAAA	TACAGAAACC	. 3780
ATCACTTTC	A ACAGTGCGAA	GACCACGCGG	CGACCTTGAA	AACCCTTTCG	CGTTCGGCCC	3840
TGAACTGCC	T TAACCCCGGA	GGCACCCTCG	TGGTGAAGTC	CTACGGTTAC	GCCGACCGCA	3900
ATAGTGAGG	A CGTAGTCACC	GCTCTTGCCA	GAAAATTTGT	CAGAGTGTCT	GCAGCGAGGC	3960
CAGAGTGCG	T CTCAAGCAAT	ACAGAAATGT	ACCTGATTTT	CCGACAACTA	GACAACAGCC	4020
GCACACGAC	A ATTCACCCCG	CATCATTTGA	ATTGTGTGAT	TTCGTCCGTG	TACGAGGGTA	4080
CAAGAGACG	G AGTTGGAGCC	GCACCGTCGT	ACCGTACTAA	AAGGGAGAAC	ATTGCTGATT	4140
GTCAAGAGG	A AGCAGTTGTC	AATGCAGCCA	ATCCACTGGG	CAGACCAGGA	GAAGGAGTCT	4200
CCCCTCCC	T CTATAAACGT	TECCCGAACA	GTTTCACCGA	TTCAGCCACA	GAGACAGGTA	4260

CCGCAAAACT GACTGTGTGC CAAGGAAAGA AAGTGATCCA CGCGGTTGGC CCTGATTTCC	1320
•	4380
	4440
	4500
	4560
GAACIGAIGC GGACGIAACC ATCIACIGCE TGGATAAGAA GTGGATGA	4620
CEGTECTCCA ACTIAAGAG TOTGTAACTG AGCTGAAGGA TGAGGATATG GTGATGAG	4680
ACGAGITAGI ATGGATCCAT CCGGACAGIT GCCTGATGGG 71 GTT GCTGATGGG 71 GTT GCCTGATGGG 71 GTT GCCTGATGGG 71 GTT GCCTGATGGG 71 GTT GCCTGATGGG 71 GT	4740
CAAAAGGAAA GIIGIAIICG IACIIIGAAG GCACCAAAII COCACCAAAA	4800
TGGCGGAGAT AAAGGICCIG IICCCAAAIG ACCAGGAAAG CACCAACAA CIGIGAGAA	4860
ACATATTGGG GGAGACCATG GAAGCAATCC GCGAAAATCC GCGAAAATCC GCGAAATCC GCAAATCC GCAAATC	
CGTCTAGCCC GCCAAAAACG CTGCCGTGCC TCTGTATGTA TGCCATGACG CCAGAAAGGG	4920
TCCACAGACT CAGAAGCAAT AACGTCAAAG AAGTTACAGT ATGCTCCTCC ACCCCCCTTC	4980
CAAAGTACAA AATCAAGAAT GTTCAGAAGG TTCAGTGCAC AAAAGTAGTC CTGTTTAACC	5040
CGCATACCCC CGCATTCGTT CCCGCCCGTA AGTACATAGA AGCACCAGAA CAGCCTGCAG	5100
CTCCGCCTGC ACAGGCCGAG GAGGCCCCCG GAGTTGTAGC GACACCAACA CCACCTGCAG	5160
CTGATAACAC CTCGCTTGAT GTCACGGACA TCTCACTGGA CATGGAAGAC AGTAGCGAAG	5220
GCTCACTCTT TTCGAGCTTT AGCGGATCGG ACAACTACCG AAGGCAGGTG GTGGTGGCTG	5280
ACGTCCATGC CGTCCAAGAG CCTGCCCCTG TTCCACCGCC AAGGCTAAAG AAGATGGCCC	5340
GCCTGGCAGC GGCAAGAATG CAGGAAGAGC CAACTCCACC GGCAAGCACC AGCTCTGCGG	5400
ACGAGTCCCT TCACCTTTCT TTTGATGGGG TATCTATATC CTTCGGATCC CTTTTCGACG	5460
GAGAGATGGC CCGCTTGGCA GCGGCACAAC CCCCGGCAAG TACATGCCCT ACGGATGTGC	5520
CTATGTCTTT CGGATCGTTT TCCGACGGAG AGATTGAGGA GTTGAGCCGC AGAGTAACCG	5580
AGTCGGAGCC CGTCCTGTTT GGGTCATTTG AACCGGGCGA AGTGAACTCA ATTATATCGT	5640
CCCGATCAGC CGTATCTTTT CCACCACGCA AGCAGAGACG TAGACGCAGG AGCAGGAGGA	5700
CCGAATACTG TCTAACCGGG GTAGGTGGGT ACATATTTTC GACGGACACA GGCCCTGGGC	5760
ACTTGCAAAA GAAGTCCGTT CTGCAGAACC AGCTTACAGA ACCGACCTTG GAGCGCAATG	5820
TTCTGGAAAG AATCTACGCC CCGGTGCTCG ACACGTCGAA AGAGGAACAG CTCAAACTCA	5880
GGTACCAGAT GATGCCCACC GAAGCCAACA AAAGCAGGTA CCAGTCTCGA AAAGTAGAAA	5940
ACCAGAAAGC CATAACCACT GAGCGACTGC TTTCAGGGCT ACGACTGTAT AACTCTGCCA	6000
CAGATCAGCC AGAATGCTAT AAGATCACCT ACCCGAAACC ATCGTATTCC AGCAGTGTAC	6060
CAGCGAACTA CTCTGACCCA AAGTTTGCTG TAGCTGTTTG TAACAACTAT CTGCATGAGA	6120
ATTACCCGAC GGTAGCATCT TATCAGATCA CCGACGAGTA CGATGCTTAC TTGGATATGG	6180
TAGACGGGAC AGTCGCTTGC CTAGATACTG CAACTTTTTG CCCCGCCAAG CTTAGAAGTT	6240
ACCCGAAAAG ACACGAGTAT AGAGCCCCAA ACATCCGCAG TGCGGTTCCA TCAGCGATGC	6300

AGAACACGTT	GCAAAACGTG	CTCATTGCCG	CGACTAAAAG	AAACTGCAAC	GTCACACAAA	6360
TGCGTGAACT	GCCAACACTG	GACTCAGCGA	CATTCAACGT	TGAATGCTTT	CGAAAATATG	6420
CATGCAATGA	CGAGTATTGG	GAGGAGTTTG	CCCGAAAGCC	AATTAGGATC	ACTACTGAGT	6480
TCGTTACCGC	ATACGTGGCC	AGACTGAAAG	GCCCTAAGGC	CGCCGCACTG	TTCGCAAAGA	6540
CGCATAATTT	GGTCCCATTG	CAAGAAGTGC	CTATGGATAG	ATTCGTCATG	GACATGAAAA	6600
GAGACGTGAA	AGTTACACCT	GGCACGAAAC	ACACAGAAGA	AAGACCGAAA	GTACAAGTGA	6660
TACAAGCCGC	AGAACCCCTG	GCGACCGCTT	ACCTATGCGG	GATCCACCGG	GAGTTAGTGC	6720
GCAGGCTTAC	AGCCGTTTTG	CTACCCAACA	TTCACACGCT	CTTTGACATG	TCGGCGGAGG	6780
ACTITGATGC	AATCATAGCA	GAACACTTCA	AGCAAGGTGA	CCCGGTACTG	GAGACGGATA.	6840
TCGCCTCGTT	CGACAAAAGC	CAAGACGACG	CTATGGCGTT	AACCGGCCTG	ATGATCTTGG	6900
AAGACCTGGG	TGTGGACCAA	CCACTACTCG	ACTTGATCGA	GTGCGCCTTT	GGAGAAATAT	6960
CATCCACCCA	TCTGCCCACG	GGTACCCGTT	TCAAATTCGG	GGCGATGATG	AAATCCGGAA	7020
TGTTCCTCAC	GCTCTTTGTC	AACACAGTTC	TGAATGTCGT	TATCGCCAGC	AGAGTATTGG	7080
AGGAGCGGCT	TAAAACGTCC	AAATGTGCAG	CATTTATCGG	CGACGACAAC	ATTATACACG	7140
GAGTAGTATC	TGACAAAGAA	ATGGCTGAGA	GGTGTGCCAC	CTGGCTCAAC	ATGGAGGTTA	7200
AGATCATTGA	CGCAGTCATC	GGCGAGAGAC	CACCTTACTT	CTGCGGTGGA	TTCATCTTGC	7260
AAGATTCGGT	TACCTCCACA	GCGTGTCGCG	TGGCGGACCC	CTTGAAAAGG	CTGTTTAAGT	7320
TGGGTAAACC	GCTCCCAGCC	GACGATGAGC	AAGACGAAGA	CAGAAGACGC	GCTCTGCTAG	7380
ATGAAACAAA	GGCGTGGTTT	AGAGTAGGTA	TAACAGACAC	CTTAGCAGTG	GCCGTGGCAA	7440
CTCGGTATGA	GGTAGACAAC	ATCACACCTG	TCCTGCTGGC	ATTGAGAACT	TTTGCCCAGA	7500
GCAAAAGAGC	ATTTCAAGCC	ATCAGAGGGG	AAATAAAGCA	TCTCTACGGT	GGTCCTAAAT	7560
AGTCAGCATA	GTACATTTCA	TCTGACTAAT	ACCACAACAC	CACCACCATG	AATAGAGGAT	7620
TCTTTAACAT	GCTCGGCCGC	CGCCCCTTCC	CAGCCCCCAC	TGCCATGTGG	AGGCCGCGGA	7680
GAAGGAGGCA	GCCGCCCCG	ATGCCTGCCC	GCAATGGGCT	GGCTTCCCAA	ATCCAGCAAC	7740
TGACCACAGC	CGTCAGTGCC	CTAGTCATTG	GACAGGCAAC	TAGACCTCAA	ACCCCACGCC	7800
CACGCCCGCC	GCCGCGCCAG	AAGAAGCAGG	CGCCAAAGCA	ACCACCGAAG	CCGAAGAAAC	7860
CAAAAACACA	GGAGAAGAAG	AAGAAGCAAC	CTGCAAAACC	CAAACCCGGA	AAGAGACAGC	7920
GTATGGCACT	TAAGTTGGAG	GCCGACAGAC	TGTTCGACGT	CAAAAATGAG	GACGGAGATG	7980
TCATCGGGCA	CGCACTGGCC	ATGGAAGGAA	AGGTAATGAA	ACCACTCCAC	GTGAAAGGAA	8040
CTATTGACCA	CCCTGTGCTA	TCAAAGCTCA	AATTCACCAA	GTCGTCAGCA	TACGACATGG	8100
AGTTCGCACA	GTTGCCGGTC	AACATGAGAA	GTGAGGCGTT	CACCTACACC	AGTGAACACC	8160
CTGAAGGGTT	CTACAACTGG	CACCACGGAG	CGGTGCAGTA	TAGTGGAGGC	AGATTTACCA	8220
TCCCCGCGG	AGTAGGAGGC	AGAGGAGACA	GTGGTCGTCC	GATTATGGAT	AACTCAGGCC	8280
GGGTTGTCGC	GATAGTCCTC	GGAGGGGCTG	ATGAGGGAAC	AAGAACCGCC	CTTTCGGTCG	8340

TCACCTGGAA TAGCAAAGGG AAGACAATCA AGACAACCCC GGAAGGGACA GAAGAGTGGT	8400
	8460
GCCCGCCCAC ATGCTACACC CGCGAACCAT CCAGAGCTCT CGACATCCTC GAAGAGAACG	8520
TGAACCACGA GGCCTACGAC ACCCTGCTCA ACGCCATATT GCGGTGCGGA TCGTCCGGCA	8580
GAAGTAAAAG AAGCGTCACT GACGACTTTA CCTTGACCAG CCCGTACTTG GGCACATGCT	8640
CGTACTGTCA CCATACTGAA CCGTGCTTTA GCCCGATTAA GATCGAGCAG GTCTGGGATG	8700
AAGCGGACGA CAACACCATA CGCATACAGA CTTCCGCCCA GTTTGGATAC GACCAAAGCG	8760
GAGCAGCAAG CTCAAATAAG TACCGCTACA TGTCGCTCGA GCAGGATCAT ACTGTCAAAG	8820
AAGGCACCAT GGATGACATC AAGATCAGCA CCTCAGGACC GTGTAGAAGG CTTAGCTACA	8880
AAGGATACTT TCTCCTCGCG AAGTGTCCTC CAGGGGACAG CGTAACGGTT AGCATAGCGA	8940
GTAGCAACTC AGCAACGTCA TGCACAATGG CCCGCAAGAT AAAACCAAAA TTCGTGGGAC	9000
GGGAAAAATA TGACCTACCT CCCGTTCACG GTAAGAAGAT TCCTTGCACA GTGTACGACC	9060
GTCTGAAAGA AACAACCGCC GGCTACATCA CTATGCACAG GCCGGGACCG CATGCCTATA	9120
CATCCTATCT GGAGGAATCA TCAGGGAAAG TTTACGCGAA GCCACCATCC GGGAAGAACA	9180
TTACGTACGA GTGCAAGTGC GGCGATTACA AGACCGGAAC CGTTACGACC CGTACCGAAA	9240
TCACGGGCTG CACCGCCATC AAGCAGTGCG TCGCCTATAA GAGCGACCAA ACGAAGTGGG	9300
TCTTCAACTC GCCGGACTCG ATCAGACACG CCGACCACAC GGCCCAAGGG AAATTGCATT	9360
TGCCTTTCAA GCTGATCCCG AGTACCTGCA TGGTCCCTGT TGCCCACGCG CCGAACGTAG	9420
TACACGGCTT TAAACACATC AGCCTCCAAT TAGACACAGA CCATCTGACA TTGCTCACCA	9480
CCAGGAGACT AGGGGCAAAC CCGGAACCAA CCACTGAATG GATCATCGGA AACACGGTTA	9540
GAAACTTCAC CGTCGACCGA GATGGCCTGG AATACATATG GGGCAATCAC GAACCAGTAA	9600
GGGTCTATGC CCAAGAGTCT GCACCAGGAG ACCCTCACGG ATGGCCACAC GAAATAGTAC	9660
AGCATTACTA TCATCGCCAT CCTGTGTACA CCATCTTAGC CGTCGCATCA GCTGCTGTGG	9720
CGATGATGAT TGGCGTAACT GTTGCAGCAT TATGTGCCTG TAAAGCGCGC CGTGAGTGCC	· 9780
TGACGCCATA TGCCCTGGCC CCAAATGCCG TGATTCCAAC TTCGCTGGCA CTTTTGTGCT	9840
GTGTTAGGTC GGCTAATGCT GAAACATTCA CCGAGACCAT GAGTTACTTA TGGTCGAACA	9900
GCCAGCCGTT CTTCTGGGTC CAGCTGTGTA TACCTCTGGC CGCTGTCGTC GTTCTAATGC	9960
GCTGTTGCTC ATGCTGCCTG CCTTTTTTAG TGGTTGCCGG CGCCTACCTG GCGAAGGTAG	10020
ACGCCTACGA ACATGCGACC ACTGTTCCAA ATGTGCCACA GATACCGTAT AAGGCACTTG	10080
TTGAAAGGGC AGGGTACGCC CCGCTCAATT TGGAGATTAC TGTCATGTCC TCGGAGGTTT	10140
TGCCTTCCAC CAACCAAGAG TACATTACCT GCAAATTCAC CACTGTGGTC CCCTCCCCTA	10200
AAGTCAGATG CTGCGGCTCC TTGGAATGTC AGCCCGCCGC TCACGCAGAC TATACCTGCA	10260
AGGTCTTTGG AGGGGTGTAC CCCTTCATGT GGGGAGGAGC ACAATGTTTT TGCGACAGTG	1032
ASSESSED FATEACTERS ECETACETCS AATTGTCAGT AGATTGCGCG ACTGACCACG	1038

CGCAGGCGAT TAAGGTGCAT ACTGCCGCGA TGAAAGTAGG ACTGCGTATA GTGTACG	GGGA 10440
ACACTACCAG TTTCCTAGAT GTGTACGTGA ACGGAGTCAC ACCAGGAACG TCTAAAC	GACC 10500
TGAAAGTCAT AGCTGGACCA ATTTCAGCAT TGTTTACACC ATTCGATCAC AAGGTCG	STTA 10560
TCAATCGCGG CCTGGTGTAC AACTATGACT TTCCGGAATA CGGAGCGATG AAACCAG	GGAG 10620
CGTTTGGAGA CATTCAAGCT ACCTCCTTGA CTAGCAAAGA CCTCATCGCC AGCACAG	ACA 10680
TTAGGCTACT CAAGCCTTCC GCCAAGAACG TGCATGTCCC GTACACGCAG GCCGCAT	TCTG 10740
GATTCGAGAT GTGGAAAAAC AACTCAGGCC GCCCACTGCA GGAAACCGCC CCTTTTG	GGT 10800
GCAAGATTGC AGTCAATCCG CTTCGAGCGG TGGACTGCTC AJACGGGAAC ATTCCCA	ATTT 10860
CTATTGACAT CCCGAACGCT GCCTTTATCA GGACATCAGA TGCACCACTG GTCTCAA	ACAG 10920
TCAAATGTGA TGTCAGTGAG TGCACTTATT CAGCGGACTT CGGAGGGATG GCTACCC	TGC 10980
AGTATGTATC CGACCGCGAA GGACAATGCC CTGTACATTC GCATTCGAGC ACAGCAA	ACCC 11040
TCCAAGAGTC GACAGTTCAT GTCCTGGAGA AAGGAGCGGT GACAGTACAC TTCAGCA	ACCG 11100
CGAGCCCACA GGCGAACTTC ATTGTATCGC TGTGTGGTAA GAAGACAACA TGCAATG	CAG 11160
AATGCAAACC ACCAGCTGAT CATATCGTGA GCACCCCGCA CAAAAATGAC CAAGAAT	TCC 11220
AAGCCGCCAT CTCAAAAACT TCATGGAGTT GGCTGTTTGC CCTTTTCGGC GGCGCCT	CGT 11280
CGCTATTAAT TATAGGACTT ATGATTTTTG CTTGCAGCAT GATGCTGACT AGCACAC	GAA 11340
GATGACCGCT ACGCCCCAAT GACCCGACCA GCAAAACTCG ATGTACTTCC GAGGAAC	TGA 11400
TGTGCATAAT GCATCAGGCT GGTATATTAG ATCCCCGCTT ACCGCGGGCA ATATAGC	AAC 11460
ACCAAAACTC GACGTATTTC CGAGGAAGCG CAGTGCATAA TGCTGCGCAG TGTTGCC	AAA 11520
TAATCACTAT ATTAACCATT TATTCAGCGG ACGCCAAAAC TCAATGTATT TCTGAGG	AAG 11580
CATGGTGCAT AATGCCATGC AGCGTCTGCA TAACTTTTTA TTATTTCTTT TATTAAT	CAA 11640
CAAAATTITG TTTTTAACAT TTC	11663

10

25

That Which Is Claimed Is:


- 1. A recombinant DNA comprising a cDNA coding for an infectious South African Arbovirus No. 86 (S.A.AR86) virus RNA transcript, and a heterologous promoter positioned upstream from said cDNA and operatively associated therewith.
- 2. The recombinant DNA according to Claim 1, wherein said cDNA is selected from the group consisting of (i) cDNA having the sequence given herein as SEQ ID NO.: 1, (ii) cDNA having the same protein coding sequence as the cDNA given herein as SEQ ID NO.: 1, and (iii) cDNA according to (i) or (ii) above, and further comprising at least one attenuating mutation in said cDNA.
 - 3. The recombinant DNA according to Claim 1, wherein said cDNA has the sequence given herein as SEQ ID NO.: 1.
- 4. The recombinant DNA according to Claim 1, further comprising at least one attenuating mutation in said cDNA clone.
 - 5. The recombinant DNA according to Claim 1, further comprising at least two attenuating mutations in said cDNA clone.
- The recombinant DNA according to Claim 5, wherein each of said attenuating mutations are in the region selected from the group consisting of the nsP1 coding region, E2 coding region, and nsP2 coding region.
 - 7. The recombinant DNA according to Claim 1, further comprising at least one attenuating mutation selected from the group consisting of codons at nsP1 amino acid position 538 which specify an attenuating amino acid, codons at E2 amino acid position 304 which specify an attenuating amino acid, codons at E2 amino acid position 314 which specify an attenuating amino acid, codons at E2 amino acid position 372 which specify an attenuating amino acid,

codons at E2 amino acid position 376 which specify an attenuating amino acid, codons at nsP2 amino acid position 96 which specify an attenuating amino acid, codons at nsP2 amino acid position 372 which specify an attenuating amino acid, codons at nsP2 amino acid position 529 which specify an attenuating amino acid, codons at nsP2 amino acid position 571 which specify an attenuating amino acid, codons at nsP2 amino acid position 682 which specify an attenuating amino acid, codons at nsP2 amino acid position 804 which specify an attenuating amino acid, and codons at nsP3 amino acid position 22 which specify an attenuating amino acid.

- 10 8. The recombinant DNA according to Claim 7, wherein said attenuating mutation comprises a substitution mutation.
 - 9. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for isoleucine at nsP1 amino acid 538.
- 10. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for threonine at E2 amino acid 304.
 - 11. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for lysine at E2 amino acid 314.
 - 12. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for valine at E2 amino acid 372.
- 20 13. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for alanine at E2 amino acid 376.
 - 14. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for glycine at nsP2 amino acid 96.

- 15. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for valine at nsP2 amino acid 372.
- 16. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for leucine at nsP2 amino acid 529.
- 5 17. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for asparagine at nsP2 amino acid 571.
 - 18. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for arginine at nsP2 amino acid 682.
- 19. The recombinant DNA according to Claim 8, wherein said10 substitution mutation codes for arginine at nsP2 amino acid 804.
 - 20. The recombinant DNA according to Claim 8, wherein said substitution mutation codes for arginine at nsP3 amino acid 22.
 - 21. The recombinant DNA according to Claim 1 further comprising at least one silent mutation.
- 15 22. The recombinant DNA according to Claim 21, wherein said silent mutation is located at a position selected from the group consisting of nucleotide 215, nucleotide 3863, nucleotide 5984 and nucleotide 9113.
- The recombinant DNA according to Claim 1, wherein not more than eight nucleotides are positioned between said promoter and said cDNA
 clone.
 - 24. The recombinant DNA according to Claim 23, wherein said promoter is selected from the group consisting of T3 promoters, T7 promoters, and SP6 promoters.

- 25. The recombinant DNA according to Claim 1, wherein said recombinant DNA comprises a plasmid, and wherein said recombinant DNA further comprises a unique restriction site positioned downstream from said cDNA clone.
- 5 26. An infectious RNA transcript encoded by a cDNA according to Claim 1.
 - 27. Infectious attenuated viral particles containing an RNA transcript of Claim 26.
- 28. A pharmaceutical composition comprising an effective immunogenic amount of an infectious attenuated virus according to Claim 27 in combination with a pharmaceutically acceptable carrier.



SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No. PCT/US96/07457

4 05		·
A. CL IPC(6)	ASSIFICATION OF SUBJECT MATTER	
US CL	:A61K 39/12; C07H 21/02; C12N 7/04, 15/40, 15/86 :Please See Extra Sheet.	
According	to International Patent Classification (IPC) or to both national classification and IPC	•
	ELDS SEARCHED	· · · · · · · · · · · · · · · · · · ·
Minimum	documentation searched (classification system followed by classification symbols)	
	435/235.1, 236, 320.1, 69.1; 424/93.1, 93.2, 93.6; 536/23,1, 23.72	•
Document	ation searched other than minimum documentation to the extent that such documents are include	ed in the fields searched
Electronic	data base consulted during the international search (name of data-base and, where practicable	e, search terms used)
APS, Di	alog, Medline, Biotech, Biosis	
Search 1	terms: alphavirus, Sindbis, arbovirus, vector, attenuated	
C. DOC	CUMENTS CONSIDERED TO BE RELEVANT	
Category*		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Α	US 5,217,879 A (HUANG et al.) 08 JUNE 1993, columns 1-	1-28
	2.	
^	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	
Α	VOS et al. Infectious RNA transcripts derived from full-length	1-28
	DNA copies of the genomic RNAs of cowpea mosaic virus.	
	Virology, 1988, Vol. 165, pages 33-41, especially page 34.	
A	VAN DED MEDE TO A COMMENT	
^	VAN DER WERF et al. Synthesis of infectious poliovirus	1-28
	RNA by purified T7 RNA polymerase. Proceedings of the	
İ	National Academy of Sciences. April 1986, Vol. 83, pages	
.]	2330-2334, especially page 2331.	
	·	
-	_	
		·
1		
		1
Furthe	er documents are listed in the continuation of Box C. See patent family annex.	
	ment defining the agreed state of the agreement in the agreement defining the agreement defining the agreement agreement defining the agreement de	tion but cited to understand the
10 6-	c of particular relevance	
	er document published on or after the international filing date "X" document of particular relevance; the considered povel or cannot be consider	e claimed invention cannot be
cited	ment which may throw doubts on priority claim(s) or which is when the document is taken alone to establish the publication date of another citation or other	
spec	an reason (as specified) Y document of particular relevance; the considered to involve an inventive	step when the document is
m can	being obvious to a person skilled in the	documents, such combination
docu the p	ment published prior to the international filing date but later than . & document member of the same patent (family
ate of the a	ctual completion of the international search Date of mailing of the international sear	rch report
17 ATICHE	/ 0 5 CED 1086	
17 AUGUS	1 1770]/
ame and ma	tiling address of the ISA/US Authorized officer : ////	1:1: 10
Box PCT	prof Patents and Trademarks	All'in gr
Washington, acsimile No.	700 000 000	()
	(703) 305-3230 Telephode No. (703) 308-0196	$V \perp$

INTERNATIONAL SEARCH REPORT

International application No. PCT/US96/07457

A. CLASSIFICATION OF SUBJECT MATTER: US CL :

435/235.1, 236, 320.1, 69.1; 424/93.1, 93.2, 93.6; 536/23,1, 23.72

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

BLACK BORDERS

IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

FADED TEXT OR DRAWING

BLURRED OR ILLEGIBLE TEXT OR DRAWING

SKEWED/SLANTED IMAGES

COLOR OR BLACK AND WHITE PHOTOGRAPHS

GRAY SCALE DOCUMENTS

LINES OR MARKS ON ORIGINAL DOCUMENT

REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

-					
	•				
				•	
•	••				
	•	•			
				,	
•			•		
					,
					,